

Article type : Case Report or Case Series

**Title:** Denosumab-related osteonecrosis of the jaw following non-surgical periodontal therapy: A case report.

**Authors:** Márcio Diniz-Freitas<sup>1</sup>; Javier Fernández-Feijoo<sup>1</sup>; Pedro Diz<sup>1</sup>; Xiana Pousa<sup>2</sup>; Jacobo Limeres<sup>1</sup>.

<sup>1</sup> Medical-Surgical Dentistry Research Group (OMEQUI), Health Research Institute of Santiago de Compostela (IDIS), University of Santiago de Compostela (USC), Santiago de Compostela, Spain.

<sup>2</sup> Periodontology Unit, School of Medicine and Dentistry, University of Santiago de Compostela (USC), Santiago de Compostela, Spain.

**Funding:** No specific funding was obtained for this study.

**Conflicts of interest:** The authors have stated explicitly that there are no conflicts of interest in connection with this article.

**Correspondence:**

Márcio Diniz Freitas

School of Medicine and Dentistry

University of Santiago de Compostela

Calle Entreríos s/n

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/jcpe.12882

This article is protected by copyright. All rights reserved.

15782 Santiago de Compostela, Spain

Email: marcio.diniz@usc.es

## ABSTRACT

**Introduction:** Osteonecrosis of the jaw associated with bisphosphonates is currently called medication-related osteonecrosis of the jaw (MRONJ), given that in addition to bisphosphonates, jaw osteonecrosis has been related to the administration of other antiresorptive and antiangiogenic drugs, such as denosumab, sunitinib, bevacizumab and ipilimumab.

**Case Presentation:** A 77-year-old patient with osteoporosis treated with subcutaneous injections of denosumab at an interval of 6 months is presented. The patient developed MRONJ after receiving a non-surgical periodontal therapy. Although the MRONJ was initially classified as a stage I lesion in this patient, cone beam computed tomography images confirmed the presence of a significant osteolytic lesion. Treatment consisted of the administration of chlorhexidine mouthwash and systemic doxycycline, exodontia of the involved teeth, sequestrectomy and complete surgical debridement of the necrotic bone.

**Conclusion:** To our knowledge, this is the first case reported in the literature of MRONJ following nonsurgical periodontal therapy in a patient with osteoporosis treated with denosumab. The risk of MRONJ development after a periodontal procedure and how to prevent this complication are still unknown.

**Keywords:** Medication-related osteonecrosis of the jaw, Denosumab.

## CLINICAL RELEVANCE

**Scientific rationale for the study:** The early diagnosis of medication-related osteonecrosis of the jaw (MRONJ) can prevent or reduce its morbidity. Although the prevalence of MRONJ is low among patients diagnosed with osteoporosis, the number of cases is increasing in view of the widespread use of receptor activator of nuclear factor kappa-B ligand (RANKL) inhibitor therapies.

**Principal findings:** The first case report on MRONJ following nonsurgical periodontal therapy in an osteoporotic patient treated with denosumab.

**Practical implications:** Dentists should be aware of the possible diagnosis of MRONJ in patients receiving or who have received antiresorptive or antiangiogenic drugs, even after dental procedures that are considered low-risk.

## INTRODUCTION

Bisphosphonate (BP)-associated osteonecrosis of the jaw (ONJ) was initially described in 2003 (Marx, 2003) and represents a complication of antiresorptive therapy especially prevalent in patients with cancer (Khan et al., 2015). Currently, it is denoted as medication-related osteonecrosis of the jaw (MRONJ), and its definition corresponds to an area of exposed bone or bone that can be probed through an intraoral or extraoral fistula that has persisted for longer than 8 weeks in patients who have received antiresorptive or antiangiogenic drugs, such as BPs, denosumab, sunitinib, bevacizumab and ipilimumab, who have neither undergone radiotherapy nor presented evidence of metastasis in this anatomic region (Ruggiero et al., 2014).

Denosumab consists of a fully human monoclonal antibody (IgG2) that mimics the action of osteoprotegerin, the physiological inhibitor of RANKL, binding to its analogue receptor, receptor activator of nuclear factor kappa-B (RANK), on the surfaces of osteoclasts (Charopoulos et al., 2011).

Denosumab acts as a regulator of bone remodeling, improving the structure, density and strength of the bone and significantly decreasing the incidence of fractures in patients with osteoporosis (Silva-Fernández et al., 2013). Denosumab also reduces the incidence of skeletal-related events in patients with cancer (Lipton, 2016) and is prescribed for the treatment of hypercalcemia of malignancy in these patients (Thosani & Hu, 2015).

Recent studies suggest that the rate of MRONJ in patients with cancer treated with RANKL inhibitors at a standard dose (120 mg of denosumab every 4 weeks) is similar to the rate found in patients receiving zoledronate (Henry et al., 2011, Van den Wyngaert et al., 2011); and it is significantly higher than the rate in patients with osteoporosis/osteopenia receiving denosumab (Papapoulos et al., 2012), generally at a dose of 60 mg subcutaneously every 6 months (O'Halloran et al., 2014). Only 2.1% of all MRONJ cases diagnosed in patients with osteoporosis have been related to the administration of denosumab; therefore, the incidence of denosumab-related ONJ is considered very low (Aljohani et al, 2017). In 2016, Bagán et al. found only 15 cases in the literature, to which they added a series of 10 new patients (Bagán et al., 2016).

A case of MRONJ following a nonsurgical periodontal therapy in a patient with osteoporosis undergoing denosumab administration is described in the present study.

## **CASE REPORT**

A 77-year-old woman was referred to the Special Care Dentistry Department of the University of Santiago de Compostela (Spain) in December 2016. Three months earlier, she had received basic periodontal therapy (scaling and root planing) for the treatment of chronic periodontitis (Figure 1). Since then, an asymptomatic bone exposure had been detected at the level of the lower incisors, which showed increased mobility (Figure 2). It should be noted that the patient's history included the discontinuous administration of BPs over the last 10 years, including ibandronic acid (150 mg

monthly) for discontinuous periods of 6–12 months, the last in 2013. In 2016, the patient started receiving biannual injections of 60 mg of denosumab subcutaneously (Prolia®, Amgen S.A., Barcelona, Spain), with a total of 2 injections; the last injection was administered in August 2016, and prednisone was concomitantly administered for 2 months.

The intraoral radiographies revealed the presence of a mixed radiolucent/radiopaque lesion suggesting bone sequestrations located between the lower lateral incisors (Figure 3). Cone beam computed tomography images confirmed the presence of an osteolytic lesion that destroyed the buccal and lingual bone plates that covered the incisor dental group and the lingual plate at the level of the lower left canine (Figure 4). The diagnosis of stage I MRONJ was established based on the clinical-radiological findings (exposed and necrotic bone in asymptomatic patients without evidence of infection).

Laboratory tests revealed C-terminal cross-linked telopeptide of type I collagen and N-terminal propeptides of type 1 procollagen serum levels fell within the postmenopausal reference range (223 pg/ml and 39.2 ng/ml, respectively). Serum levels of alkaline phosphatase, hydroxyvitamin D and calcium were normal.

Surgical treatment of the lesion was planned, and antimicrobial treatment with a 0.2% chlorhexidine mouthwash (twice per day) and doxycycline (200 mg/day) was established from 7 days prior to the surgery until 3 weeks after the surgical procedure. Extraction of the teeth involved in the sequestration and complete surgical debridement of the necrotic bone until reaching clinically vital bone margins were performed (Figure 5), followed by a primary closure. The patient is asymptomatic, without any evidence of bone exposure or continuity solution of the covering mucosa after 6 months of clinical and radiological follow-up (Figure 6).

## DISCUSSION AND CONCLUSION

Taking into account that marketing and clinical use of denosumab were approved in 2010, many patients diagnosed with MRONJ were previously treated with BPs (Aljohani et al., 2017). It is likely that inhibition of bone resorption by BPs was still present when our patient started denosumab and osteonecrosis developed, which makes it difficult to establish the role of each of these drugs in the development of the MRONJ, as has been suggested in previous series (Bagán et al., 2016). However, a time frame longer than 12 months between the discontinuation of BPs and the onset of denosumab treatment, as in the present case, is sufficient for the bone to recover its metabolic activity, and, consequently, to be able to attribute the appearance of MRONJ to denosumab (Pichardo & van Merkesteyn, 2016).

Our patient developed MRONJ 4 years after the suspension of BP treatment and after receiving only 2 injections of denosumab. The treatment duration represents one of the main risk factors for the development of BP-related ONJ (Palaska et al., 2009). Recently, it has been estimated that a period of 6.0 and 2.2 years of oral alendronate and intravenous zoledronate therapy, respectively, is required for 50% of patients to develop MRONJ (Fung et al., 2017). In contrast, cases of denosumab-related ONJ tend to occur early. The mean number of doses of denosumab administered to patients with osteoporosis from the series published by Bagán et al. (Bagán et al., 2016) was 3, but cases of MRONJ after the administration of a single dose of denosumab without a history of prior consumption of BPs have been described (Pichardo et al, 2013), confirming that no clear association between the dose or duration of denosumab treatment and the development of MRONJ has been confirmed to date (Yoshimura et al, 2017).

Most cases of MRONJ appear as a consequence of invasive dental procedures, mainly dental extractions (Diniz-Freitas & Limeres Posse, 2016; Bagán et al., 2016). Experts have noted that the risk of MRONJ after a surgical periodontal procedure that requires bone exposure and manipulation is comparable to the risk associated with dental extractions (Ruggiero et al., 2014). In contrast,

nonsurgical periodontal procedures should not be considered “high-risk” because these procedures do not cause evident trauma to the supporting tissues. Although the actual risk of developing MRONJ from these procedures is still unknown (Ruggiero et al., 2014), some authors have insinuated that MRONJ can be caused by lesser traumas in patients treated with denosumab than in patients receiving BPs (O’Halloran et al., 2014).

It has been suggested that periodontal disease is a significant triggering factor for the development of MRONJ (Yamazaki et al., 2012; McGowan et al., 2017). This possibility is of special importance in patients who receive denosumab, given in a published series, most cases of ONJ developed spontaneously, without any known triggering event like dentoalveolar surgery, ill-fitting dentures or local trauma (Yarom et al. 2017). Anaerobic periodontopathogenic bacteria have been detected in necrotic bone in MRONJ and in perilesional areas that are clinically healthy, including *Porphyromonas gingivalis*, *Treponema denticola* and *Tannerella forsythia* (Hallmer, 2017). However, their role in the etiopathogenesis of MRONJ has not yet been clarified.

In the present case, nonsurgical periodontal treatment (scaling and root planing), which was likely the triggering procedure, was conducted only 1 month after the last injection of denosumab. There is no evidence in patients undergoing treatment with BPs that protocols of temporary drug interruption (i.e., drug holidays) can prevent the development of MRONJ secondary to invasive dental procedures (Diniz-Freitas & Limeres-Posse, 2016; Hasegawa et al., 2017). This is probably due to the fact that BPs, once incorporated into the bone matrix, remain united to the bone for years (e.g., the terminal half life of alendronate is approximately 10 years). In contrast to BPs, denosumab does not bind to hydroxyapatite; therefore, the effect of denosumab on bone remodeling, although more efficient, is less durable (Russel et al., 2008; Fukumoto & Matsumoto, 2017), and is mostly ineffectual within 6 months of treatment cessation (Ruggiero et al., 2014). Accordingly, some authors have proposed interruption protocols for denosumab of 6 months prior to a surgical procedure to facilitate bone regeneration (Epstein et al., 2013, Robinson & Scully, 2014).

There is no high-quality evidence on the efficacy of antibiotic prophylaxis for patients treated with antiresorptive drugs who undergo dental extractions. However, it appears that antibiotics combined with other perioperative measures, such as osteotomy and primary wound closure, exert a certain preventive effect (Gaudin et al., 2015; Diniz-Freitas and Limeres-Posse, 2016). The efficacy of antibiotic prophylaxis in preventing medication-related osteonecrosis of the jaw (MRONJ) before undergoing periodontal therapy for patients treated with receptor activator of nuclear factor kappa-B ligand (RANKL) inhibitors is, however, still unknown. The treatment modalities for MRONJ are typically classified as pharmacological and surgical. The management of stage I MRONJ includes the prescription of antimicrobial mouthwashes, clinical follow-up on a quarterly basis, patient education and the review of the indications for antiresorptive therapy (Ruggiero et al., 2014). However, the staging proposed by the American Association of Oral and Maxillofacial Surgeons (USA) underestimates the extent of the lesions, and computed tomography is required to obtain an accurate definition of the limits of the involved bone (Bedogni et al., 2014). In the present case, the advanced dental mobility and the presence of bone sequestration led us to choose a less conservative therapy in an attempt to completely remove the necrotic bone; otherwise, the risk of recurrence or progression of the disease would remain (Carlson & Basile, 2009; Carlson, 2014). In this regard, although conservative treatment can stop the progression of the disease and fight superinfection, better results have been obtained with surgical treatment; therefore, an aggressive surgical approach has been proposed as the first option in the treatment of MRONJ (El-Rabbany et al., 2017; Hayashida et al., 2017), even in the early stages of the condition (Ristow et al., 2015).

In summary, to our knowledge, this is the first case reported in the literature of MRONJ following nonsurgical periodontal therapy in a patient with osteoporosis treated with denosumab. Unfortunately, to date there have been no evidence-based protocols aimed at preventing MRONJ in

patients with periodontitis who receive denosumab, even though its pharmacokinetics could justify its suppression 6 months before starting periodontal therapy. Dentists should be aware of the development of MRONJ related to new therapeutic modalities of osteoporosis (RANKL inhibitors) including denosumab. The risk of appearance of these lesions following a periodontal procedure and how to prevent them are still unknown.

## References

- Aljohani, S., Fliefel, R., Ihbe, J., Kühnisch, J., Ehrenfeld, M. & Otto, S. (2017) What is the effect of anti-resorptive drugs (ARDs) on the development of medication-related osteonecrosis of the jaw (MRONJ) in osteoporosis patients: A systematic review. *J Craniomaxillofac Surg* (In press). doi: 10.1016/j.jcms.2017.05.028.
- Bagan, J., Peydró, A., Calvo, J., Leopoldo, M., Jiménez, Y., Bagan, L. (2016) Medication-related osteonecrosis of the jaw associated with bisphosphonates and denosumab in osteoporosis. *Oral Dis* **22**, 324-329. doi: 10.1111/odi.12447.
- Bedogni, A., Fedele, S., Bedogni, G., Scoletta, M., Favia, G., Colella, G., Agrillo, A., Bettini, G., Di Fede, O., Oteri, G., Fusco, V., Gabriele, M., Ottolenghi, L., Valsecchi, S., Porter, S., Petruzzi, M., Arduino, P., D'Amato, S., Ungari, C., Fung Polly, P.L., Saia, G. & Campisi, G. (2014) Staging of osteonecrosis of the jaw requires computed tomography for accurate definition of the extent of bony disease. *Br J Oral Maxillofac Surg* **52**, 603-608. doi: 10.1016/j.bjoms.2014.04.009.
- Carlson, E.R. & Basile, J.D. (2009) The role of surgical resection in the management of bisphosphonate-related osteonecrosis of the jaws. *J Oral Maxillofac Surg* **67**, 85–95.

Carlson E.R. (2014) Management of antiresorptive osteonecrosis of the jaws with primary surgical resection. *J Oral Maxillofac Surg* **72**, 655–657. doi: 10.1016/j.joms.2013.12.007. doi: 10.1016/j.joms.2009.01.006.

Charopoulos, I., Orme, S., Giannoudis, P.V. (2011) The role and efficacy of denosumab in the treatment of osteoporosis: an update. *Expert Opin Drug Saf* **10**, 205–217. doi: 10.1517/14740338.2010.516249.

Cremers, S.C., Pillai, G. & Papapoulos, S.E. (2005) Pharmacokinetics-pharmacodynamics of bisphosphonates. *Clin Pharmacokinet* **44**:551–570.

Diniz-Freitas, M. & Limeres-Posse, J. (2016) Prevention of medication-related osteonecrosis of the jaws secondary to tooth extractions. A systematic review. *Med Oral Patol Oral Cir Bucal* **21**: e250-e259. doi:10.4317/medoral.20963.

El-Rabbany, M., Sgro, A., Lam, D.K., Shah, P.S. & Azarpazhooh, A. (2017) Effectiveness of treatments for medication related osteonecrosis of the jaw: A systematic review and meta-analysis. *J Am Dent Assoc* **148**, 584-594.e2. doi: 10.1016/j.adaj.2017.04.002.

Epstein, M.S., Ephros, H.D. & Epstein, J.B. (2013) Review of current literature and implications of RANKL inhibitors for oral health care providers. *Oral Surg Oral Med Oral Pathol Oral Radiol* **116**, e437-e442. doi: 10.1016/j.oooo.2012.01.046.

Fukumoto, S. & Matsumoto, T. (2017) Recent advances in the management of osteoporosis. *F1000Res* **6**,625. doi: 10.12688/f1000research.10682.1.

Fung, P., Bedogni, G., Bedogni, A., Petrie, A., Porter, S., Campisi, G., Bagan, J., Fusco V., Saia, G., Acham, S., Musto, P., Petrucci, M.T., Diz, P., Colella, G., Mignogna, M.D., Pentenero, M., Arduino, P., Lodi, G., Maiorana, C., Manfredi, M., Hallberg, P., Wadelius, M., Takaoka, K., Leung, Y.Y., Bonacina, R., Schiødt, M., Lakatos, P., Taylor, T., De Riu, G., Favini, G., Rogers,

S.N., Pirmohamed, M., Nicoletti, P.; GENVABO Consortium & Fedele, S. (2017) Time to onset of bisphosphonate-related osteonecrosis of the jaws: a multicentre retrospective cohort study. *Oral Dis* **23**, 477-483. doi: 10.1111/odi.12632.

Gaudin, E., Seidel, L., Bacevic, M., Rompen, E. & Lambert, F. (2015) Occurrence and risk indicators of medication-related osteonecrosis of the jaw after dental extraction: a systematic review and meta-analysis. *J Clin Periodontol* **42**, 922-32. doi: 10.1111/jcpe.12455

Hasegawa, T., Kawakita, A., Ueda, N., Funahara, R., Tachibana, A., Kobayashi, M., Kondou, E., Takeda, D., Kojima, Y., Sato, S., Yanamoto, S., Komatsubara, H., Umeda, M., Kirita, T., Kurita, H., Shibuya, Y., Komori, T & Japanese Study Group Cooperative Dentistry with Medicine (JCDM). (2017) A multicenter retrospective study of the risk factors associated with medication-related osteonecrosis of the jaw after tooth extraction in patients receiving oral bisphosphonate therapy: can primary wound closure and a drug holiday really prevent MRONJ? *Osteoporos Int* **28**, 2465-2473. doi: 10.1007/s00198-017-4063-7.

Hayashida, S., Soutome, S., Yanamoto, S., Fujita, S., Hasegawa, T., Komori, T., Kojima, Y., Miyamoto, H., Shibuya, Y., Ueda, N., Kirita, T., Nakahara, H., Shinohara, M. & Umeda, M. (2017) Evaluation of the Treatment Strategies for Medication-Related Osteonecrosis of the Jaws (MRONJ) and the Factors Affecting Treatment Outcome: A Multicenter Retrospective Study with Propensity Score Matching Analysis. *J Bone Miner Res* **32**, 2022-2029. doi: 10.1002/jbmr.3191.

Hallmer, F., Bjørnland, T., Andersson, G., Becktor, J.P., Kristoffersen, A.K. & Enersen, M. (2017) Bacterial diversity in medication-related osteonecrosis of the jaw. *Oral Surg Oral Med Oral Pathol Oral Radiol* **123**, 436-444. doi: 10.1016/j.oooo.2016.11.011

Henry, D.H., Costa, L., Goldwasser, F., Hirsh, V., Hungria, V., Prausova, J., Scagliotti, G.V., Sleeboom, H., Spencer, A., Vadhan-Raj, S., von Moos, R., Willenbacher, W., Woll, P.J., Wang, J., Jiang, Q.,

Jun, S., Dansey, R. & Yeh, H. (2011) Randomized, double-blind study of denosumab versus zoledronic acid in the treatment of bone metastases in patients with advanced cancer (excluding breast and prostate cancer) or multiple myeloma. *J Clin Oncol* **29**, 1125-1132. doi: 10.1200/JCO.2010.31.3304.

Khan, A.A., Morrison, A., Hanley, D.A., Felsenberg, D., McCauley, L.K, O'Ryan, F., Reid, I.R., Ruggiero, S.L., Taguchi, A., Tetradis, S., Watts, N.B., Brandi, M.L., Peters, E., Guise, T., Eastell, R., Cheung, A.M., Morin, S.N., Masri, B., Cooper, C., Morgan, S.L., Obermayer-Pietsch, B., Langdahl, B.L., Al Dabagh, R., Davison, K.S., Kendler, D.L., Sándor, G.K., Josse, R.G., Bhandari, M., El Rabbany, M., Pierroz, D.D., Sulimani, R., Saunders, D.P., Brown, J.P., Compston, J & International Task Force on Osteonecrosis of the Jaw. (2015) Diagnosis and management of osteonecrosis of the jaw: a systematic review and international consensus. *J Bone Miner Res* **30**, 3-23. doi: 10.1002/jbmr.2405.

Kendler, D.L., Roux, C., Benhamou, C.L., Brown, J.P., Lillestol, M., Siddhanti S., Man, H.S., San Martin, J. & Bone, H.G. (2010) Effects of denosumab on bone mineral density and bone turnover in postmenopausal women transitioning from alendronate therapy. *J Bone Miner Res* **25**, 72-81. doi: 10.1359/jbmr.090716.

Lipton, A., Fizazi, K., Stopeck, A.T., Henry, D.H., Smith, M.R., Shore, N., Martin, M., Vadhan-Raj, S., Brown, J.E., Richardson, G.E., Saad, F., Yardley, D.A., Zhou, K., Balakumaran, A. & Braun, A. (2016) Effect of denosumab versus zoledronic acid in preventing skeletal-related events in patients with bone metastases by baseline characteristics. *Eur J Cancer* **53**: 75-83. doi:10.1016/j.ejca.2015.09.011.

Marx, R.E. (2003) Pamidronate (Aredia) and zoledronate (Zometa) induced avascular necrosis of the jaws: a growing epidemic. *J Oral Maxillofac Surg* **61**, 1115-1117. doi: dx.doi.org/10.1016/S0278-2391(03)00720-1.

MacGowan, K., McGowan, T. & Ivanovski, S. (2017) Risk factors for medication-related osteonecrosis of the jaws: A systematic review. *Oral Dis* (In press). doi: 10.1111/odi.12708

O'Halloran, M, Boyd, N.M. & Smith, A. (2014) Denosumab and osteonecrosis of the jaws - the pharmacology, pathogenesis and a report of two cases. *Aust Dent J* **59**, 516-519. doi: 10.1111/adj.12217.

Palaska, P.K., Cartsos, V. & Zavras, A.I. (2009) Bisphosphonates and time to osteonecrosis development. *Oncologist* **14**, 1154-1166. doi: 10.1634/theoncologist.2009-0115.

Papapoulos, S., Chapurlat, R., Libanati, C., Brandi, M.L., Brown, J.P., Czerwiński, E., Krieg, M.A., Man, Z., Mellström, D., Radominski, S.C., Reginster, J.Y., Resch, H., Román Ivorra, J.A., Roux, C., Vittinghoff, E., Austin, M., Daizadeh, N., Bradley, M.N., Grauer, A., Cummings S.R. & Bone, H.G. (2012) Five years of denosumab exposure in women with postmenopausal osteoporosis: results from the first two years of the FREEDOM extension. *J Bone Miner Res* **27**:694-701. doi: 10.1002/jbmr.1479.

Pichardo, S.E., Kuypers, S.C. & van Merkesteyn, J.P. (2013) Denosumab osteonecrosis of the mandible: a new entity? A case report. *J Craniomaxillofac Surg* **41**, e65-9. doi: 10.1016/j.jcms.2012.10.014.

Pichardo, S.E. & van Merkesteyn, J.P. (2016) Evaluation of a surgical treatment of denosumab-related osteonecrosis of the jaws. *Oral Surg Oral Med Oral Pathol Oral Radiol* **122**, 272-278. doi: 10.1016/j.oooo.2016.03.008.

Ramírez, L., López-Pintor, R.M., Casañas, E., Arriba, L.D. & Hernández, G. (2015) New Non-Bisphosphonate Drugs that Produce Osteonecrosis of the Jaws. *Oral Health Prev Dent* **13**, 385-393. doi: 10.3290/j.ohpd.a34055.

Ristow, O., Otto, S., Troeltzsch, M., Hohlweg-Majert, B. & Pautke, C. (2015) Treatment perspectives for medication-related osteonecrosis of the jaw (MRONJ). *J Craniomaxillofac Surg* **43**, 290-293. doi: 10.1016/j.jcms.2014.11.014.

Robinson, A. & Scully, C. (2014) Pharmacology: New therapies and challenges. *Br Dent J* **217**, 258-259. doi: 10.1038/sj.bdj.2014.811.

Ruggiero, S.L., Dodson, T.B., Fantasia, J., Goodday, R., Aghaloo, T., Mehrotra, B., O'Ryan, F. & American Association of Oral and Maxillofacial Surgeons. (2014) American Association of Oral and Maxillofacial Surgeons position paper on medication-related osteonecrosis of the jaw--2014 update. *J Oral Maxillofac Surg* **72**, 1938-1956. 10.1016/j.joms.2014.04.031.

Russell, R.G., Watts, N.B., Ebetino, F.H. & Rogers, M.J. (2008) Mechanisms of action of bisphosphonates: similarities and differences and their potential influence on clinical efficacy. *Osteoporos Int* **19**, 733-759. doi: 10.1007/s00198-007-0540-8.

Silva-Fernández, L., Rosario, M.P., Martínez-López, J.A., Carmona, L. & Loza, E. (2013) Denosumab for the treatment of osteoporosis: a systematic literature review. *Reumatol Clin* **9**, 42-52. doi: 10.1016/j.reuma.2012.06.007

Thosani, S. & Hu, M.I. (2015) Denosumab: a new agent in the management of hypercalcemia of malignancy. *Future Oncol* **11**, 2865-71. doi: 10.2217/fon.15.232.

Van den Wyngaert, T., Wouters, K., Huizing, M.T. & Vermorken, J.B. (2011) RANK ligand inhibition in bone metastatic cancer and risk of osteonecrosis of the jaw (ONJ): non bis in idem? *Support Care Cancer* **19**, 2035-2040. doi: 10.1007/s00520-010-1061-0.

Yamazaki, T., Yamori, M., Ishizaki, T., Asai, K., Goto, K., Takahashi, K., Nakayama, T. & Bessho, K. (2012) Increased incidence of osteonecrosis of the jaw after tooth extraction in patients

treated with bisphosphonates: a cohort study. *Int J Oral Maxillofac Surg* **41**, 1397-1403. doi: 10.1016/j.ijom.2012.06.020.

Yarom, N., Lazarovici, T.S., Whitefield, S., Weissman, T., Wasserzug, O. & Yahalom, R. (2018) Rapid onset of osteonecrosis of the jaw in patients switching from bisphosphonates to denosumab. *Oral Surg Oral Med Oral Pathol Oral Radiol.* **125**, 27-30. doi: 10.1016/j.oooo.2017.09.014.

Yoshimura, H., Ohba, S., Yoshida, H., Saito, K., Inui, K., Yasui, R., Ichikawa, D., Aiki, M., Kobayashi, J., Matsuda, S., Imamura, Y. & Sano, K. (2017) Denosumab-related osteonecrosis of the jaw in a patient with bone metastases of prostate cancer: A case report and literature review. *Oncol Lett* **14**, 127-136. doi: 10.3892/ol.2017.6121.

#### Figure Legends

**Figure 1.** Periodontal chart at first appointment.

**Figure 2.** Initial clinical image showing exposed bone in the buccal and lingual zones of the anterior mandible, without signs of infection.

**Figure 3.** Intraoral radiographs showing the presence of periradicular radiolucent areas.

**Figure 4.** Cone beam computed tomography (CBCT) images showing the extent of the osteolytic process affecting the buccal and lingual cortical plates.


**Figure 5.** Surgical treatment consisting of extractions of the involved teeth, sequestrectomy and a complete surgical debridement of the necrotic bone until reaching the clinically vital margins, followed by primary closure.

**Figure 6.** Clinical and radiological aspect at 6 months after surgical treatment.

## Periodontal Chart



Chart #:  
Name:  
Examiner:  
Date:



The style and appearance of this chart is a trademark of the Florida Probe Corporation-Copyright © 1996-2010, All Rights Reserved - Gainesville, Florida, USA  
Printed on 12/12/2017 16:52:07 from YOUR-152F56135B



